

## EFFECTS OF CSPN MODELS ON PNE SHELL MODELING

B. Armsdorfer<sup>1</sup>, S. Kimeswenger<sup>1</sup>, and T. Rauch<sup>2</sup>

birgit.armsdorfer@uibk.ac.at, stefan.kimeswenger@uibk.ac.at, rauch@astro.uni-tuebingen.de

**Modeling planetary nebulae (PNe) shells using different ionizing spectra for a hot central star (CSPN) of evolved PNe, we found that especially a blackbody model leads to wrong nebular diagnostics and abundances.**

The aim of our investigations is to combine modeling of stellar atmospheres and of planetary nebulae shells to obtain self consistent descriptions of the nebulae. For this purpose, spatially resolved roundish nebulae with well modeled central stars were selected (Rauch et al. 1999). As a test sample, the nebula NGC 2438 was selected, and the photo-ionization code Cloudy (Ferland 1996) was used for the modeling. To tune nebular parameters like density profile and filling factor, we used one of the new NLTE stellar model atmosphere fluxes with H-Ni element coverage by Rauch (2001) with  $T_{\text{eff}}=110$  kK and  $\log g=7$  (cgs). The other geometrical parameters were derived from narrow-band images and long-slit spectra. This model was then used to calculate four different sets of artificial narrow-band surface brightness distributions in various species (H I, He I, He II, O III, N II, S II). The sources of illumination were [a] the new NLTE model atmosphere by Rauch as mentioned above, [b] a H-Ca NLTE model (Rauch 1997), [c] a NLTE atmosphere containing only H-He, and [d] a blackbody. The stellar fluxes were calibrated to have the same visual magnitude,  $T_{\text{eff}}$  and  $\log g$ . While  $H\alpha$  and  $H\beta$  are nearly unaffected by the different illumination sources, other lines change both in strength and spatial profile. For the low ionized species, the blackbody overestimates the lines by 30% ([S II]) to 45% ([N II]) – therefore, also the ratio  $H\alpha$ /[S II] vs.  $H\alpha$ /[N II] used for PNe identification diagrams changes up to 15%. For the [O III] lines, the blackbody underestimates the real flux by 30%. He I<sub>4471</sub> is underestimated by 20%, while He II<sub>4686</sub>, which also shows the strongest spatial change, is overestimated by 270% with the blackbody model (see Fig. 1). Models [b] and [c] underestimate the flux by about 20%. All diagnostic diagrams are therefore affected. This leads typi-




Fig. 1. Results of our PNe shell modeling from the line structure output of Cloudy for He II<sub>4686</sub>, the lines are normalized to the integrated  $H\beta$  flux. The blackbody model [d] overestimates the flux, the changes from model [a] to model [c] are also visible.

cally to an overestimation of the stellar temperature and nitrogen abundance. A blackbody model for the stellar atmosphere thus leads to wrong nebular parameters and should not be used anymore. This work will be discussed more detailed in a forthcoming paper.

This project was supported by the BMBWK, Sektion VIII/A/5, and by a DLR grant 50 OR97055. Support from the organizers to participate in the conference is gratefully acknowledged.

### REFERENCES

- Ferland, G. J. 1996, Hazy, a Brief Introduction to Cloudy 90.05, Univ. Kentucky, Department of Physics and Astronomy, Internal Report
- Rauch, T. 1997, A&A, 320, 237
- Rauch, T., Köppen, J., Napiwotzki, R., & Werner, K. 1999, A&A, 347, 169
- Rauch, T. 2001, this issue

<sup>1</sup>Institut für Astrophysik, Universität Innsbruck, Austria.

<sup>2</sup>Institut für Astronomie und Astrophysik, Universität Tübingen, Germany.